

Title page

Depression and anxiety as predictors of mortality among heart failure patients: systematic review and meta-analysis

I. Sokoreli^{1,*}, J.J.G. de Vries¹, S.C. Pauws¹, E.W. Steyerberg²

(1) Philips Research - Healthcare, High Tech Campus 34, 5656 AE Eindhoven, The Netherlands; (2) Department of Public Health, Centre for Medical Decision Making, Erasmus MC, Rotterdam, the Netherlands

* Corresponding author. Tel.: +310646945879.
E-mail address: joanna.sokoreli@philips.com (I. Sokoreli).

Abstract

Aims: Several studies suggest that psychological factors are associated with negative outcomes and in particular higher mortality rates among Heart Failure (HF) patients. We aimed to evaluate the effect sizes of depression and anxiety on all-cause mortality in HF patients.

Methods and Results: We conducted a systematic review according to the PRISMA methodology. We searched for studies on depression or anxiety effects on all-cause mortality among HF patients published up to June 2015. A number of 26 and 6 articles met inclusion criteria for depression (total 80627 patients) and anxiety (total 17214 patients) respectively. The effect estimates were pooled using random-effect meta-analysis.

Depression has significant and moderately heterogeneous effect on all-cause mortality (HR = 1.57; 95%CI 1.30 – 1.89, $P < 0.001$); adjustment for confounders led to a similar effect estimate (HR = 1.40; 95%CI 1.22 – 1.60; $p < 0.001$). Larger studies and higher study prevalence of depression were associated with smaller effect size. The effect of anxiety on mortality outcome was small and not conclusive given the low number of studies ($n=6$) (HR = 1.02; 95% CI 1.00 – 1.04, $P < 0.05$).

Conclusions: This systematic review and meta-analysis suggests that depression is an important and independent predictor of all-cause mortality among HF patients, while anxiety does not appear to have a strong effect. Further research is recommended towards the detection and treatment of depression.

Keywords: heart failure, depression, anxiety, mortality, systematic review, meta-analysis.

Introduction

Heart Failure (HF) is defined as a clinical syndrome in which patients have typical symptoms such as breathlessness, ankle swelling, and fatigue and signs such as elevated jugular venous pressure, pulmonary crackles, and displaced apex beat, resulting from an abnormality of cardiac structure or function [1]. Approximately 1–2% of the adult population in developed countries has HF, with the prevalence rising to $\geq 10\%$ among persons 70 years of age or older [2]. HF is one of the most common causes of hospital readmission and mortality.

Psychological factors such as depression or anxiety are often reported with high prevalence and strong association with negative outcomes in patients with cardiovascular disease [3]. Many studies have reported high rates of depression among HF patients. A prior systematic review and meta-analysis published by Rutledge in 2006 [4] reported an overall aggregated depression prevalence rate of 21.6% among HF patients, while individual study prevalence estimates ranged from 9% to 60%. Moreover, in 2005 Konstam [5] reported that approximately 40% of HF patients may suffer from major anxiety, and overall anxiety levels are 60% higher than levels seen in the healthy population.

Depression has been linked to increased risk of negative outcomes, such as rehospitalization and mortality among HF patients. According to a previous meta-analysis, the aggregated risk estimate derived from 8 studies suggested a greater than 2-fold risk of death and secondary events for HF patients with heightened depressive symptoms or a depressive disorder [4]. A similar analysis was also published by Fan [6] in 2014 on 9 prospective studies, who reported a pooled Hazard Ratio of 1.51 for patients with depression compared to patients without depression. In both cases the result was strongly heterogeneous but no further analysis, such as meta-regression, was performed to examine the sources of this heterogeneity. On the other hand, there is, to the best of our knowledge, no meta-analysis published about the prevalence of anxiety among HF patients and the effect of anxiety on mortality outcome. Even though anxiety is usually correlated with depression, it has not extensively been studied among patients with HF.

Our aim is to provide an updated systematic review of prospective or retrospective studies and a meta-analysis of the effect of depression and the effect of anxiety on mortality among HF patients. To reach this objective, we searched

extensively for available studies investigating the impact of depression and anxiety on mortality of HF patients. Within these studies, we identified also the reported prevalence of depression or anxiety among HF patients.

Methods

Search strategy and selection criteria

This systematic review and meta-analysis were conducted according to the guidelines introduced in the Preferred Reporting Items for Systematic reviews and Meta-analysis (the PRISMA Statement) [7]. The 27 checklist items of the PRISMA methodology followed are given in Appendix A. Three electronic databases (MEDLINE, BIOSIS and EMBASE) were searched for studies that investigated the relationship between depression or anxiety and mortality among Heart Failure (HF) patients. No publication time restriction was applied. All papers written in English and published before the 25th of June 2015 were included. Selected journals as well as the references of full-text papers were also hand-searched, when necessary, in order to identify studies that meet the inclusion criteria.

The database search string was created according to the PICO model (P, population/patient; I, intervention/indicator; C, comparator/control; and O, outcome). For the “P” in PICO the “HEART FAILURE” keyword was included. For the “I”, the following keywords: “DEPRESS? OR STRESS OR ANXIETY OR PSYCHOLOG?”. For the “C”, no particular terms were used in our case. For “O”, we used the following keywords: “MORTALITY OR DEATH”. The complete query as used for the databases search is given in Appendix B.

Study selection

In our analysis, several inclusion and exclusion criteria were defined. All studies that met those criteria were included. The inclusion criteria were articles presenting studies focusing on the association between depression or anxiety and mortality in a HF adult population. All mortality outcomes such as all-cause or cardiac related mortality were included and studies focusing on inpatient, outpatient or both care settings were taken into account. On the other hand, publications analyzing data that had already been used before for the same purpose, studies introducing no quantitative assessment of the impact of depression or anxiety on the outcome or analyzing the use of antidepressants as primary focus were excluded from our analysis.

Review process and data collection

All titles and abstracts of studies identified by the electronic and hand search were screened by the reviewer (IS) to identify those meeting the inclusion/exclusion criteria. Then, all the selected full texts were screened independently by two reviewers (IS, GJdV) to identify which articles should be included in the systematic review. Any disagreement between the reviewers was resolved by a third reviewer (SP). For each of the selected articles the reviewers extracted data about author, year of publication, follow-up period, outcome variable, location, study design, study population (size/ type), prevalence of depression or anxiety, assessment method of the psychological parameter, other parameters, statistical method and results.

Mendeley 1.13.8 software was used for organizing and managing of the articles.

Data analysis

All studies were categorized according to the psychological factor investigated (depression or anxiety). Information was extracted according to whether the analysis was adjusted for confounders such as age, gender, and clinical severity. For both groups the association between depression or anxiety and mortality was reported by collecting information of the hazard ratios/odds ratios, 95%CI and/or p-values.

Random-effects meta-analysis was applied to combine the results. We decided to pool not only the adjusted effect but also the unadjusted effects in order to avoid the bias of the different adjustments. For the few cases where Odds Ratios were reported, they were converted [8] into Hazard Ratios in order to be comparable with the other Hazard Ratios. In studies where results were presented for several periods of follow-up we selected the longest follow-up period to avoid bias of including multiple results on the same patient data.

Studies collected in our analysis were different with respect to patient population, locations and depression or anxiety assessment methods. The random-effects method allows for heterogeneity by assuming that the effects being estimated in the different studies are not identical, but follow a normal distribution. Heterogeneity across the studies was quantified by the I^2 statistic [9]. The I^2 statistic summarizes the fraction of the variation across studies due to heterogeneity relative to chance. Random-effect meta-regression was used in an attempt to explain between-

study heterogeneity and identify possible sources of bias. Meta-regression is a method to quantify the association between the estimated effect of depression and different study characteristics.

Meta analyses were presented in the form of forest plots created with the metafor package for R statistics version 3.0.3 (The R Foundation for Statistical Computing).

Results

Search result

A total of 906 potentially relevant articles was identified from the electronic search and 5 from the hand search. After removing the duplicates and reviewing the titles and abstracts we ended up with 62 articles for a full text review. From these, 35 more articles were excluded, leaving 27 articles for the systematic review (Fig. 1).

→ Fig. 1

Characteristics of the selected studies

Depression and mortality

Among the identified studies, 26 reported on the effect of depression. The prevalence of depression varied from 10 to 79% in the identified literature studies (Table 1). The unadjusted effect of depression is presented in Table 1a, while the effect of depression after adjusting for several confounders in Table 1b. The most common confounders, used in more than 10 studies, were age, gender, NYHA class and (left ventricular) ejection fraction.

There were various techniques used among the studies to assess depression levels. We included all studies assessing for clinically significant depression. The most common scale used was the Beck Depression Inventory (BDI) [10], followed by the Patient Health Questionnaire (PHQ) [11].

→ Table 1

The pooled hazard ratio for the unadjusted effect of depression on mortality was strongly significant across 15 studies (HR = 1.57; 95%CI 1.30 – 1.89; $p < 0.001$). The pooled estimation was strongly heterogeneous as reflected by the I^2 statistic ($I^2 = 94\%$, heterogeneity $p < 0.001$). The pooled adjusted Hazard Ratio was also significant (HR = 1.40; 95%CI 1.22 – 1.60; $p < 0.001$) and again heterogeneous (heterogeneity $p < 0.001$; $I^2 = 97\%$, Fig. 2)

→ Fig. 2

A random-effect meta-regression was performed to understand the sources of the higher than 90% observed heterogeneity between the studies. The potential study-level covariates analyzed were the study characteristics introduced in Table 1. There was no association found between heterogeneity and the depression assessment method, the adjusted or univariate analysis, the location where the study was conducted, the inpatient or outpatient predictive period, the year of the study, the type of the study and the follow-up period. On the other hand, significant heterogeneity was associated with the total population size (smaller effect in larger studies $p < 0.01$) and the prevalence of the depression in the study (smaller effect for prevalence $>29\%$; $p < 0.01$, Table 2).

→ Table 2

Anxiety and mortality

Only 6 studies analyzing the effect of anxiety on mortality among HF patients were identified with a prevalence of anxiety varying from 9 to 53% (Table 3). Table 3a shows the unadjusted effects reported in the studies and Table 3b the reported effects on mortality after adjusting for a group of confounders. Age, NYHA class and (left ventricular) ejection fraction were the most common confounders in the identified studies.

→ Table 3

There was no evidence found for anxiety as an independent predictor of mortality. The pooled hazard ratio for the unadjusted effect of anxiety on mortality, which was based on 2 studies, was 1.02 (95% CI 1.00 – 1.04; $p = 0.24$, heterogeneity $p = 0.38$; $I^2 = 0\%$). The pooled hazard ratio for the adjusted effect of anxiety on mortality could be based on 5 studies and was identical (HR = 1.02; 95% CI 1.00 – 1.04; $p = 0.09$) and reasonably homogenous (heterogeneity $p = 0.97$; $I^2 = 0\%$, Fig. 3).

→ Fig. 3

Discussion

This systematic review was conducted according to the PRISMA guidelines to assess the evidence on the effect of depression (26 studies) and anxiety (6 studies) on all-cause mortality outcome among Heart Failure (HF) patients.

<Key results: 1.6 for depression but very heterogeneous across studies; no effect for anxiety>.

In contrast to other reviews, our study was not limited on follow-up duration or only in prospective studies reporting adjusted effects of the two parameters. We reviewed all studies published quantifying the effect of depression or anxiety.

The prevalence of depression varied among the 26 different studies with an average of approximately 29% ranging from 10 to 79%. The meta-analysis showed that the unadjusted risk of death among HF patients facing depression was 1.57 times higher than the risk among HF patients without depression and the pooled estimate of the adjusted Hazard Ratio was 1.40. In both univariate and adjusted analysis, strong heterogeneity among the studies was found. Our findings are more conservative than previous reviews published [4, 6]. Rutledge et al reported a 2.10 higher adjusted risk of mortality and secondary events based on 8 studies and Fun et al reported a pooled adjusted Hazard Ratio of 1.51 based on 9 studies, both with substantial heterogeneity. From our attempt to explain heterogeneity we found that the effect of depression is weaker in larger studies; this suggests publication bias: small studies were published if they found relatively large effect estimates, while small studies with modest effect estimates were not. The weaker effect in studies with higher prevalence of depression may relate to the use of different cut-offs on an underlying, latent, scale for depression. If a more liberal cut-off was used, those labeled as depressed actually were milder than with a more strict definition of depression.

Our results for anxiety do not have the same weight as the results with respect to depression since anxiety was less studied in the literature. Anxiety had a similar prevalence to depression among the six identified studies (average 29%, range 9 - 45%), but patients with anxiety had no increased risk of death compared to those without anxiety. However, since anxiety is usually correlated with other factors such as depression, further research of anxiety as a covariate to other factors is recommended.

One limitation of our study is related to the variation in follow-up times. Follow up times varied from 30 days to a number of years; furthermore, there were studies covering different follow up periods but in these cases we always selected the longest follow up. Further analysis such as subgroup analysis would be recommended to investigate the effect variation in different follow-up periods, however limited information in some of the literature publications is restrictive towards this direction.

Moreover, we focused only on mortality. Nevertheless, there is evidence that depression and anxiety are also associated to other adverse events such as readmission. Further investigation is needed also towards this direction.

One limitation of the meta-regression is that even though we tried to cover a broad selection of study-level covariates there are more that might also be related to the heterogeneity. Further research on different factors' interactions would be recommended.

The "gold standard" test of causality of a putative risk factor is a randomized clinical trial. Such a trial minimizes concerns about confounders [12–14]. To the best of our knowledge, there is no randomized clinical trial conducted for depression among a HF population. Based on our findings we strongly recommend such a trial in order to evaluate the causality of depression.

Finally, according to our findings from the meta-regression, depression should not be underestimated in clinical practice within HF population groups where prevalence is low. Furthermore, based on our overall findings on the effect of depression, we recommend further research on the recognition and management of depression in clinical practice which might improve patient outcomes. Further analysis such as subgroup analysis and interventional studies are required for stronger evidence towards this direction.

Acknowledgements

The authors would like to thank John Dielis from E-Science & Information Management Department, Philips, Eindhoven for contributing in querying the databases.

Funding and Conflict of interest

IS, JJGV and SCP are employed by Philips Research. EWS has no conflict of interest to declare.

References

1. McMurray JJ V, Adamopoulos S, Anker SD, et al. (2012) ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012. *Eur J Heart Fail* 14:803–869. doi: 10.1093/eurjhf/hfs105
2. Mosterd A, Hoes AW (2007) Clinical epidemiology of heart failure. *Heart* 93:1137–1146. doi: 10.1136/hrt.2003.025270
3. Ladwig K-H, Lederbogen F, Albus C, et al. (2014) Position paper on the importance of psychosocial factors in cardiology: Update 2013. *Ger Med Sci* 12:Doc09. doi: 10.3205/000194
4. Rutledge T, Reis V a., Linke SE, et al. (2006) Depression in Heart Failure. A Meta-Analytic Review of Prevalence, Intervention Effects, and Associations With Clinical Outcomes. *J Am Coll Cardiol* 48:1527–1537. doi: 10.1016/j.jacc.2006.06.055
5. Konstam V, Moser DK, De Jong MJ (2005) Depression and anxiety in heart failure. *J Card Fail* 11:455–463. doi: 10.1016/j.cardfail.2005.03.006
6. Fan H, Yu W, Zhang Q, et al. (2014) Depression after heart failure and risk of cardiovascular and all-cause mortality: A meta-analysis. *Prev Med (Baltim)* 63:36–42. doi: 10.1016/j.ypmed.2014.03.007
7. Liberati A, Altman DG, Tetzlaff J, et al. (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *Ann Intern Med*. doi: 10.1371/journal.pmed.1000100
8. Higgins JPT, Altman D, Gøtzsche P, et al. (2011) The Cochrane Collaboration’s tool for assessing risk of bias in randomized trials. doi: 10.1002/9780470712184
9. Higgins JPT, Thompson SG, Deeks JJ, Altman DG (2003) Measuring inconsistency in meta-analyses. *BMJ Br Med J* 327:557–560. doi: 10.1136/bmj.327.7414.557
10. Zich JM, Attkisson CC, Greenfield TK (1990) Screening for depression in primary care clinics: the CES-D and the BDI. *Int J Psychiatry Med* 20:259–277.
11. Kroenke K, Spitzer RL, Williams JBW (2001) The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med* 16:606–613. doi: 10.1046/j.1525-1497.2001.016009606.x
12. Ioannidis JP a. (2005) Contradicted and Initially Stronger Effects in Highly Cited Clinical Research. *JAMA J Am Med Assoc* 294:218–228. doi: 10.1001/jama.294.2.218
13. Evans I, Thornton H, Chalmers I (2008) Testing treatments - Better research for better healthcare. *Chinese J Evidence-Based Med*. doi: 10.1136/bmj.332.7556.1516
14. Mushlin AI, Ghomrawi H (2010) Health care reform and the need for comparative-effectiveness research. *N Engl J Med* 362:e6. doi: 10.1056/NEJMp0912651
15. Adams J, Kuchibhatla M, Christopher EJ, et al. (2012) Association of Depression and Survival in Patients with Chronic Heart Failure over 12 Years. *Psychosomatics* 53:339–346. doi: 10.1016/j.psym.2011.12.002

16. Albert NM, Fonarow GC, Abraham WT, et al. (2009) Depression and Clinical Outcomes in Heart Failure: An OPTIMIZE-HF Analysis. *Am J Med* 122:366–373. doi: 10.1016/j.amjmed.2008.09.046
17. Diez-Quevedo C, Lupón J, González B, et al. (2013) Depression, antidepressants, and long-term mortality in heart failure. *Int J Cardiol* 167:1217–1225. doi: 10.1016/j.ijcard.2012.03.143
18. Faller H, Störk S, Schowalter M, et al. (2007) Depression and survival in chronic heart failure: Does gender play a role? *Eur J Heart Fail* 9:1018–1023. doi: 10.1016/j.ejheart.2007.06.011
19. Faller H, Störk S, Gelbrich G, et al. (2015) Depressive symptoms in heart failure: Independent prognostic factor or marker of functional status? *J Psychosom Res.* doi: 10.1016/j.jpsychores.2015.02.015
20. Farisa R, Purcell H, Henein MY, Coats a. JS (2002) Clinical depression is common and significantly associated with reduced survival in patients with non-ischaemic heart failure. *Eur J Heart Fail* 4:541–551. doi: 10.1016/S1388-9842(02)00101-0
21. Friedmann E, Thomas S a., Liu F, et al. (2006) Relationship of depression, anxiety, and social isolation to chronic heart failure outpatient mortality. *Am Heart J* 152:1–8. doi: 10.1016/j.ahj.2006.05.009
22. Jiang W, Alexander J, Christopher E, et al. (2001) Relationship of depression to increased risk of mortality and rehospitalization in patients with congestive heart failure. *Arch Intern Med* 161:1849–1856. doi: 10.1001/archinte.161.15.1849
23. Jiang W, Kuchibhatla M, Clary GL, et al. (2007) Relationship between depressive symptoms and long-term mortality in patients with heart failure. *Am Heart J* 154:102–108. doi: 10.1016/j.ahj.2007.03.043
24. Jünger J, Schellberg D, Müller-Tasch T, et al. (2005) Depression increasingly predicts mortality in the course of congestive heart failure. *Eur J Heart Fail* 7:261–267. doi: 10.1016/j.ejheart.2004.05.011
25. Kato N, Kinugawa K, Yao A, et al. (2009) Relationship of depressive symptoms with hospitalization and death in Japanese patients with heart failure. *J Card Fail* 15:912–919. doi: 10.1016/j.cardfail.2009.06.442
26. Lesman-Leegte I, Van Veldhuisen DJ, Hillege HL, et al. (2009) Depressive symptoms and outcomes in patients with heart failure: Data from the COACH study. *Eur J Heart Fail* 11:1202–1207. doi: 10.1093/eurjhf/hfp155
27. Moraska AR, Chamberlain AM, Shah ND, et al. (2013) Depression, healthcare utilization, and death in heart failure a community study. *Circ Heart Fail* 6:387–394. doi: 10.1161/CIRCHEARTFAILURE.112.000118
28. O'Connor CM, Abraham WT, Albert NM, et al. (2008) Predictors of mortality after discharge in patients hospitalized with heart failure: An analysis from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *Am Heart J* 156:662–673. doi: 10.1016/j.ahj.2008.04.030
29. Sullivan MD, Levy WC, Crane B a., et al. (2004) Usefulness of depression to predict time to combined end point of transplant or death for outpatients with advanced heart failure. *Am J Cardiol* 94:1577–1580. doi: 10.1016/j.amjcard.2004.08.046
30. Alhurani AS, Dekker RL, Abed M a., et al. (2014) The Association of Co-morbid Symptoms of Depression and Anxiety With All-Cause Mortality and Cardiac Rehospitalization in Patients With Heart Failure. *Psychosomatics* 56:371–80. doi: 10.1016/j.psym.2014.05.022

31. Coyne JC, Rohrbaugh MJ, Shoham V, et al. (2001) Prognostic importance of marital quality for survival of congestive heart failure. *Am J Cardiol* 88:526–529. doi: 10.1016/S0002-9149(01)01731-3
32. Cully J a, Johnson M, Moffett ML, et al. (2009) Depression and anxiety in ambulatory patients with heart failure. *Psychosomatics* 50:592–598. doi: 10.1176/appi.psy.50.6.592
33. Konstam V, Salem D, Pouleur H, et al. (1996) Baseline Quality of Life as a Predictor of Mortality and Hospitalization in 5,025 Patients With Congestive Heart Failure. *Am J Cardiol* 78:890–895. doi: 10.1016/S0002-9149(96)00463-8
34. Murberg T a, Bru E (2001) Social relationships and mortality in patients with congestive heart failure. *J Psychosom Res* 51:521–527.
35. Rollman BL, Herbeck Belnap B, Mazumdar S, et al. (2012) A positive 2-item patient health questionnaire depression screen among hospitalized heart failure patients is associated with elevated 12-month mortality. *J Card Fail* 18:238–245. doi: 10.1016/j.cardfail.2011.11.002
36. Sherwood A, Blumenthal J a, Trivedi R, et al. (2007) Relationship of depression to death or hospitalization in patients with heart failure. *Arch Intern Med* 167:367–373. doi: 10.1001/archinte.167.4.367
37. Smith ORF, Kupper N, Schiffer a. a., Denollet J (2012) Somatic Depression Predicts Mortality in Chronic Heart Failure: Can This Be Explained by Covarying Symptoms of Fatigue? *Psychosom Med* 74:459–463. doi: 10.1097/PSY.0b013e31824ef2f4
38. Van Den Broek KC, Defilippi CR, Christenson RH, et al. (2011) Predictive value of depressive symptoms and B-type natriuretic peptide for new-onset heart failure and mortality. *Am J Cardiol* 107:723–729. doi: 10.1016/j.amjcard.2010.10.055
39. Volz A, Schmid J-P, Zwahlen M, et al. (2011) Predictors of readmission and health related quality of life in patients with chronic heart failure: a comparison of different psychosocial aspects. *J Behav Med* 34:13–22. doi: 10.1007/s10865-010-9282-8
40. Zuluaga MC, Guallar-Castillón P, Rodríguez-Pascual C, et al. (2010) Mechanisms of the association between depressive symptoms and long-term mortality in heart failure. *Am Heart J* 159:231–237. doi: 10.1016/j.ahj.2009.11.011
41. Jiang W, Kuchibhatla M, Cuffe MS, et al. (2004) Prognostic value of anxiety and depression in patients with chronic heart failure. *Circulation* 110:3452–3456. doi: 10.1161/01.CIR.0000148138.25157.F9

Figure legends

Fig. 1 PRISMA flow diagram	29
Fig. 2 Meta-analysis – Forest plot calculating the effect of depression (a) unadjusted effect, (b) adjusted effect	30
Fig. 3 Meta-analysis – Forest plot calculating the effect of anxiety (a) unadjusted effect, (b) adjusted effect	31

APPENDICES

Appendix A PRISMA checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	3-4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	Appendix A
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5-6

Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5-6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Appendix B
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5-6
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	5-7
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6-7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	6-7
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ²) for each meta-analysis.	6-7
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	6-7
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	6-7
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Fig.1

Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Table 1 – Table 3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	/
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Table 1 – Table 3; Fig. 2-3
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Fig. 2-3
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	9/Table 2
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	9/Table 2
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	10-11
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	10-11
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	10-11
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	12

Appendix B Database search query

S (HEART(W)FAILURE)/TI AND ((DEPRESS? OR STRESS? OR ANXIETY OR
PSYCHOLOG?)(S)(MORTALITY OR DEATH))/TI,AB

Tables

Table 1 (a) Unadjusted/ (b) adjusted effect of depression on mortality among HF patients

(a) Unadjusted effect of depression on mortality													
Author	Year	Assessment method	Outcome	Population	Region	Study	Follow-up	Predicting period	Statistical method	HR/OR	p-value	95% CI	Prevalence of depression
Adams [15]	2012	BDI score ≥ 10	all-cause mortality	985 HF	US	Prospective cohort study	mean 1792.3 days	-	Univariate Cox proportional-hazards model	1.35	<0.001	1.15–1.57	30%
Albert [16]	2009	history of depression	all-cause mortality	48612 HF	US	OPTIMIZE-HF comprehensive registry	60-90 days	Inpatient	Univariate Cox proportional-hazards model	1.36	0.027	1.04- 1.79	11%
Diez-Quevedo[17]	2013	GDS-4 ≥ 1	all-cause mortality	1017 HF	ES	Prospective cohort study	median 5.4 years	Outpatient	Univariate Cox proportional-hazards model	1.39	0.001	1.15-1.68	42%
Faller [18]	2007	PHQ-9 - major	all-cause mortality	231 CHF	DE	Prospective cohort study	median 2.7 years	Outpatient	Univariate Cox proportional-hazards model	3.30	<0.001	1.8–6.1	13%
Faller [19]	2015	PHQ-9 score	all-cause mortality	863 HF	DE	Extended INH study	18 months	Outpatient	Univariate Cox proportional-hazards model	1.07	<0.001	1.04–1.09	-
Faris [20]	2002	ICD-10	all-cause mortality	39 HF	UK	Retrospective cohort study	mean 48 months	Outpatient	Univariate Cox proportional-hazards model	2.10	0.0005	1.4-3.2	21%
Friedmann [21]	2006	BDI-II increase of 1	all-cause mortality	231 CHF	US	PFOS cohort study	mean 23.6 months	Outpatient	Univariate Cox proportional-hazards model	2.59	0.0177	0.231-5.431	36%
Jiang [22]	2001	BDI ≥ 10	all-cause mortality	374 CHF	US	Prospective cohort study	1 year	Inpatient	Univariate logistic regression	2.26	0.04	1.04-4.91	35%
Jiang [23]	2007	BDI ≥ 10	all-cause mortality	1006 HF	US	Cohort study	mean 971 days	Inpatient	Univariate Cox proportional-hazards model	1.45	<0.001	1.19-1.77	30%
Junger [24]	2005	HADS-D > 6	all-cause mortality	209 CHF	DE	Prospective study	mean 24.8 months	-	Univariate Cox proportional-hazards model	1.09	0.0071	1.02–1.17	30%
Kato [25]	2009	CES-D ≥ 16	all-cause mortality	115 HF	JP	Prospective cohort study	median 2.1 years	Outpatient	Univariate Cox proportional-hazards model	5.51	0.004	1.75-17.39	23%
Lesman-	2009	CES-D ≥ 24	all-cause	958	NL	COACH prospective	18 months	Inpatient	Univariate Cox proportional-	1.18	0.1724	0.93 - 1.5	21%

Leegte [26]			mortality	HF		study				hazards model				
Moraska [27]	2013	PHQ-9 \geq 10	all-cause mortality	402 HF	US	Prospective cohort study	mean 1.6 years	in/outpatient	Univariate Cox proportional-hazards model	3.37	<0.001	1.97-5.75	15%	
O'connor [28]	2008	history of depression	all-cause mortality	5791 HF	US	OPTIMIZE-HF Prospective cohort study	mean 72.7 days	Inpatient	Univariate Cox proportional hazards model	1.56	0.0004	1.23-1.97	14%	
Sullivan [29]	2004	PRIME-MD interview/HDRS/ SCL-20	all-cause mortality	142 HF	US	Prospective cohort study	mean 3 years	Outpatient	Univariate Cox proportional-hazards model	1.65	0.403	0.51-5.28	29%	
(b) Adjusted effect of depression on mortality														
Author	Year	Assessment method	Outcome	Population	Region	Study	Follow-up	Predicting period	Statistical method	Other parameters	HR /OR	p-value	95% CI	Prevalence of depression
Adams [15]	2012	BDI score \geq 10	all-cause mortality	985 HF	US	Prospective Cohort study	mean 1792.3 days	-	Multivariate Cox proportional-hazards model	age, sex, race, marital status, NYHA, ischemic etiology of HF, history of CABG, diagnosis of diabetes	1.40	<0.001	1.16-1.68	30%
Albert [16]	2009	interviews/ medical records	all-cause mortality	48612 HF	US	OPTIMIZE-HF comprehensive hospital-based registry	60-90 days	Inpatient	Multivariate Cox proportional-hazards model	age, race, history of: ischemic heart disease, depression, hypertension, liver disease and diabetes, any revascularization procedure, any mechanical ventilation, discharge medication: ACE, aldosterone antagonists, digoxin and lipid-lowering agent/ discharge vital signs: SBP, DBP,HR; admission laboratory: serum sodium, discharge lab: serum creatinine	1.46	0.025	1.05-2.03	11%
Alhurani [30]	2015	PHQ-9 \geq 10	all-cause mortality	1260 HF	US	HF Health-Related QoL Collaborative Registry	12 month	Outpatient	Multivariate Cox proportional-hazards model	age, gender, ethnicity, NYHA, combined anxiety/depression	1.06	0.012	1.01-1.11	33%
Coyne [31]	2011	CES-D \geq 16	all-cause mortality	706 HF	NL	COACH study randomized control trial	18 month	Inpatient	Multivariate Cox proportional-hazards model	BNP, type D	1.01	0.066	0.10-1.03	34%
Cully [32]	2009	ICD-9 code depression	all-cause mortality	12028 HF	US	Retrospective cohort study	12 month	Outpatient	Multivariate logistic regression	age, gender, race, married, income, comorbidities, combined depression/anxiety	0.93	ns	0.71-1.15	18%
Diez-Quevedo	2013	GDS-4 \geq 1	all-cause mortality	1017 HF	ES	Prospective cohort study	median 5.4	Outpatient	Multivariate Cox proportional-hazards	Sex, age, Months since HF diagnosis, Ischemic etiology, LVEF, NYHA, DM, COPD, Peripheral	1.31	0.008	1.07-1.60	42%

[17]							years		model	vasculopathy, CrC, BMI, ACE or ARB, BB				
Faller [18]	2007	PHQ-9 major	all-cause mortality	231 CHF	DE	Prospective cohort study	median 2.7 years	Outpatient	Multivariate Cox proportional-hazards model	Age, sex, aetiology, NYHA, EF, syst./non-syst. LV dysfunction, interaction term b/w LVEF and LV dysfunction	2.40	0.008	1.3–4.6	13 %
Faller [19]	2015	PHQ-9 score	all-cause mortality	863 HF	DE	extended INH study	18 months	Outpatient	Multivariate Cox proportional-hazards model	age, sex, randomization status, NYHA, LVEF 30%, amino-terminal pro-BNP, SBP, HR, coronary artery disease, renal dysfunction, anemia, diabetes, ACE, ARB, BB, diuretics, and statins	1.04	0.017	1.01–1.07	-
Faris [20]	2002	ICD-10	all-cause mortality	396 HF	UK	Retrospective cohort study	mean 48 months	Outpatient	Multivariate Cox proportional-hazards model	demographics, social, medical history, baseline functional status and clinical severity	3.00	0.004	1.4–6.4	21%
Friedmann [21]	2006	BDI-II increase of 1	all-cause mortality	231 CHF	US	PFOS cohort study	mean 23.6 months	Outpatient	Multivariate Cox proportional-hazards model	treatment: ICD, amiodarone, afib, EF, depression score, social support amount	2.35	0.0222	2.354–4.743	36%
Jiang [22]	2001	BDI ≥ 10 and positive DIS result	all-cause mortality	374 CHF	US	Prospective cohort study	1 year	Inpatient	Multivariate logistic regression	age, LVEF, NYHA, ischemic aetiology of CHF	2.12	0.07	0.94–4.81	35%
Jiang [23]	2007	BDI ≥ 10	all-cause mortality	1006 HF	US	cohort study	mean 971 days	Inpatient	Multivariate Cox proportional-hazards model	age, LVEF, NYHA, ischemic aetiology of CHF, history of diabetes, marital status	1.40	0.003	1.12–1.74	30%
Junger [24]	2005	HADS-D > 6	all-cause mortality	209 CHF	DE	Prospective study	mean 24.8 months	-	Multivariate Cox proportional-hazards model	peakVO2, LVEF	1.08	0.02	1.01–1.15	30% (HADS-D ≥ 8)
Kato [25]	2009	CES-D ≥ 16	all-cause mortality	115 HF	JP	Prospective cohort study	median 2.1 years	Outpatient	Multivariate Cox proportional-hazards model	age, ACE, BNP	5.52	0.006	1.65–18.46	24%
Konstam [33]	1996	HRQL	all-cause mortality	3375 HF	US	Randomized clinical trial	mean 36.5 months	-	Multivariate Cox proportional-hazards model	EF, age, treatment, NYHA	1.07	0.023	1.009–1.125	-
Lesman-Leegte [26]	2009	CES-D ≥ 24	all-cause mortality	958 HF	NL	COACH Prospective study	18 months	Inpatient	Multivariate Cox proportional-hazards model	age, gender, BNP level	1.43	0.04	1.02–2.02	21%
Moraska	2013	PHQ-9 ≥	all-cause	402 HF	US	Prospective	mean	in/outpatient	Multivariate Cox	age, gender, CCI, incident vs. prevalent HF status	4.06	<0.001	2.35-	15%

[27]		10	mortality			cohort study	1.6 years		proportional-hazards model				7.01	
Murberg and Bru [34]	2001	SDS	all-cause mortality	119 CHF	NO	Prospective study	2 year	Outpatient	Multivariate Cox proportional-hazards model	depressive symptoms, NYHA, functional status and age	1.05	0.116	0.988-1.108	-
O'connor [28]	2008	history of depression	all-cause mortality	5791 HF	US	OPTIMIZE-HF prospective cohort study	mean 72.7 days	Inpatient	Multivariate stepwise Cox proportional-hazards model	SBP, age, weight, Reactive airway disease, sodium, SCr, Liver disease, Lower extremity edema, Statin at discharge, BB at discharge	1.48	0.0034	1.14-1.93	14%
Rollman [35]	2012	PHQ-2	all-cause mortality	471 HF	US	Prospective study	up to 12 months	Inpatient	Multivariate Cox proportional-hazards model	sex, age ≥ 65 , EF $\leq 30\%$, NYHA 3/4, anxiety, COPD, renal insufficiency, ACE-I or ARB, BB, Coumadin, hemoglobin < 10 , sodium < 136 , DBP, SBP	3.10	0.003	1.4-6.7	79%
Sherwood [36]	2007	BDI ≥ 10	all-cause mortality	204 HF	US	Prospective study	median 3 years	Outpatient	Multivariate Cox proportional-hazards model	NT-proBNP, antidepressant, age, HF etiology, and LVEF	1.05	0.06	1.00-1.10	46%
Smith [37]	2012	BDI score	all-cause mortality	380 CHF	NL	-	median 2.3 years	Outpatient	Multivariate Cox proportional-hazards model	male, age, LVEF, NYHA, smoking, exertion fatigue	1.41	0.02	1.05-1.88	-
van den Broek [38]	2011	CES-D ≥ 8	all-cause mortality	208 HF	NL	Prospective community based study	median 11 years	Outpatient	Multivariate Cox proportional-hazards model	age, gender, race, SBP, cholesterol, DM, BMI, smoking, reduced physical activity, CHD at baseline, LVEF, left ventricular hypertrophy, NT-proBNP	1.49	-	1.05-2.11	36%
Volz [39]	2011	HADS > 10	all-cause mortality	111 HF	CH	Prospective cohort study	mean 2.8 years	Outpatient	Multivariate Cox proportional-hazards model	LVEF, peak oxygen uptake	0.65	0.7	0.08-5.17	10%
Zuluaga [40]	2010	GDS-10 ≥ 5	all-cause mortality	433 HF	ES	Prospective study	mean 5.7 year	Outpatient	Multivariate Cox proportional-hazards model	age, gender, race, COPD, CCI, serum creatinine level, LVEF, NYHA, HF hospitalization in last year, ischemic cardiopathy, heart valve disease	1.40	< 0.01	1.05-1.86	24%

HR, Hazard Ratio; OR, Odds Ratio; BDI, Beck Depression Inventory; GDS, Geriatric Depression Scale; PHQ, Patient Health Questionnaire; HADS, Hospital Anxiety and Depression Scale; CES-D, Center for Epidemiological Studies Depression Scale; PRIME-MD, Primary Care Evaluation of Mental Health Disorders; HDRS, Hamilton Rating Scale for Depression; SCL, Symptom Checklist.; HRQL, Health Related Quality of Life; SDS, Zung Self-Rating Depression Scale; PFOS, Psychosocial Factors Outcome Study; CABG, coronary artery bypass graft; EF, Ejection Fraction; LVEF, Left Ventricular Ejection Fraction; CCI, Charlson Comorbidity Index; SBP, Systolic Blood Pressure; DBP, Diastolic Blood Pressure; HR, Heart Rate; ACE, Angiotensin-converting enzyme; ARB, Angiotensin II Receptor Blockers; BB, Beta Blockers

Table 2 Random-effect meta-regression¹

	Univariate Analysis	
	Estimated coefficient (SE)	P-value
Year	-0.0016 (0.0124)	0.8957
Assessment method		
1. BDI	0.0349 (0.1287)	0.7863
2. PHQ	0.2096 (0.1433)	0.1434
3. Other	-0.1571 (0.1117)	0.1596
Population size *	-0.0004 (0.0002)	<0.05
Region		
1. EU	-0.1119 (0.1134)	0.3241
2. US	0.04355 (0.1140)	0.7555
Follow-up period	0.0014 (0.0269)	0.9599
Statistical method		
1. Unadjusted	0.1066 (0.1159)	0.3573
2. Adjusted	reference	reference
Study type		
1. Prospective	0.1453 (0.1134)	0.2003

2. Retrospective	0.0667 (0.2217)	0.7637
3. Other	-0.1756 (0.1178)	0.1359
Depression prevalence*	-0.0108 (0.0059)	<0.1
Predicting period		
1. Inpatient	-0.1641 (0.1156)	0.1156
2. Outpatient	Reference	reference
¹ In order to estimate the unadjusted effect of each study-level factor, the studies with missing values were excluded in each case		

Table 3 (a) Unadjusted/ (b) adjusted effect of anxiety on mortality among HF patients

(a) Unadjusted effect of anxiety on mortality														
Author	Year	Assessment method	Outcome	Population	Region	Study	Follow-up	Predicting period	Statistical method	Other parameters	HR/OR	P-value	95% CI	Prevalence of anxiety
Friedmann [21]	2006	STAI	all-cause mortality	149 CHF	US	PFOS cohort study	23.6 months	Outpatient	Univariate Cox proportional-hazards model	-	1.037	0.06	0.998-1.078	45%
Jiang [41]	2004	STAI	all-cause mortality	291 CHF	US	Prospective cohort study	1 year	Inpatient	Univariate Cox proportional-hazards model	-	State-A:1.017 Trait-A:1.010	State-A:0.12 Trait-A: 0.44	State-A:0.996 – 1.039 Trait-A:0.98–1.03	29%
(b) Adjusted effect of anxiety on mortality														
Author	Year	Assessment method	Outcome	Population	Region	Study	Follow-up	Predicting period	Statistical method	Other parameters	HR/OR	P-value	95% CI	Prevalence of anxiety
Alhurani [30]	2015	BSI anxiety subscale	all-cause mortality	1260 HF	US	HF QoL Registry	12 month	Outpatient	Multivariate Cox proportional-hazards model	age, gender, ethnicity, NYHA, anxiety&depression	1.07	0.652	0.79–1.45	-
Cully [32]	2009	ICD 9 anxiety	all-cause mortality	12028 HF	US	Retrospective cohort study	12 month	Outpatient	Multivariate logistic regression	age, gender, race, married, income, comorbidities, combined depression/anxiety	1.01	ns	0.76 - 1.54	9%
Friedmann [21]	2006	STAI	all-cause mortality	149 CHF	US	PFOS cohort study	23.6 months	Outpatient	Multivariate Cox proportional-hazards model	atrial fibrillation/flutter, NYHA, and treatment group	1.03	0.12	0.989 -1.072	45%
Jiang [41]	2004	STAI ≥ 40	all-cause mortality	291 CHF	US	Prospective cohort study	1 year	Inpatient	Multivariate Cox proportional-hazards model	BDI scores, age, baseline LVEF, NYHA, and ischemic CHF origin	State-A: 1.01 Trait-A: 1.00	State-A: 0.30 Trait-A: 0.97	State-A: 0.988– 1.040 Trait-A: 0.971–1.031	-
Konstantin [33]	1996	HRQL	all-cause mortality	3375 HF	US	Randomized clinical trial	mean 36.5 months	-	Multivariate Cox proportional-hazards model	EF, age, treatment, NYHA classification, and HRQL	1.02	ns	-	-
Volz [39]	2011	HADS-A > 10	all-cause mortality	111 HF	CH	Prospective cohort study	mean 2.8 years	Outpatient	Multivariate Cox proportional-hazards model	LVEF, peak oxygen uptake	1.75	0.47	0.37–8.21	9%

Hazard Ratio(HR); Odds Ratio(OR); STAI: State-Trait Anxiety Inventor; Health Related Quality of Life (HRQL); Brief Symptom Inventory (BSI); Psychosocial Factors Outcome Study (PFOS)

Figures




Fig. 1 PRISMA flow diagram




Fig. 2 Meta-analysis – Forest plot calculating the effect of depression (a) unadjusted effect, (b) adjusted effect




Fig. 3 Meta-analysis – Forest plot calculating the effect of anxiety (a) unadjusted effect, (b) adjusted effect